STRAWBERRIES.

NAMES.

A.

G. Κοραρον

L. Fragaria.

I. *Fravola.*

G. Erdbeeren.

F. Fraises.

S. Miruedanos.

D. Aardbeeze

PARTS
Roots leave

PREPAR
None kept in

PLACE.
They are cultivated in Gardens; some grow wild in many Places.

T
Flower
ripe June

DESCRIPTION.

The Roots are small, reddish, full of Fibres, that from Runners abo shoot Fibres downward, and Leaves upward, of a middling Green, Jagg'd, whiter The Stalks rise to support the Fruit above the Leaves, which from three upon one Foot to one near the Fruit. The Empalement 2 is a very light whitish Green, set with whi divided into ten pointed Parts. The Flowers 3 have five white Leaves, in the Cente is the young Fruit: at 4 'tis cut through to shew the Embrio of the Fruit, with the Outside, with every one a Yellow Pistil, with reddish Yellow Tops; the Hives are n Summits cover'd with a Yellow Farina; the ripe Fruit has many light-coloured n on their Outside, one of each Sort is over the Fruit that produced it. The Scarle 5, the Wood 6, which is a deeper Red, and the Hoboy 7, of which there are two Sort Black, the other White: 8 shews a Fruit cut Longitudinally, 9 another Transversly is whitish.

VIRTUES.

The Leaves are used in Gargarisms, for Soreness of the Mouth &c. The Fruit at Table than in Shops.

T. Sheldrake delin.

TIMOTHY SHELDRAKE *Botanicum Medicinale (An Herbal of Medicinial Plants)* LONDON, 1759 CHICAGO BOTANIC GARDEN